Matthew H. Blake, MD

Team Physician, University of Sioux Falls Assistant Professor, Department of Orthopedic Surgery University of South Dakota School of Medicine www.MatthewBlakeMD.com



Physical Therapy Prescription – Meniscus Repair

Name:	_ Date:	
Procedure: R / L meniscus repair	Date of Surgery:	_
Frequency: 2-3 times per week for weeks		
ASE I (Waaks 0 - 6): Period of protection, decrease edema	activate quadricens	

PHASE I (Weeks 0 – 6): Period of protection, decrease edema, activate quadriceps

- Weightbearing: Partial with crutches
 - **Weeks 2-4:** advance to 50% weight-bearing in brace with crutches
 - Weeks 4-6: Progress to full weight-bearing in brace, wean off crutches
- **Hinged Knee Brace:**
 - Weeks 0-2: locked in full extension for ambulation and sleeping
 - o Weeks 2-6: unlocked (0-90°) for ambulation and removed while sleeping
- Range of Motion: AAROM → AROM 0-90°
- Therapeutic Exercises: Patellar mobs, quad/hamstring sets, heel slides, Gastroc/Soleus stretching, straight-leg raises with brace in full extension until guad strength prevents extension lag
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase II (Weeks 6 – 12)

- Weightbearing: As tolerated, unassisted
- Hinged Knee Brace: Discontinue at 6 weeks
- Range of Motion: Full: no weight-bearing with knee flexion angles >90°
- Therapeutic Exercises: Patellar mobs, quad/hamstring sets, closed chain extension exercises, hamstring curls, toe raises, balance exercises, Gastroc/Soleus stretching; lunges 0-90°, leg press 0-90°
 - Begin use of the stationary bicycle
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase III (Weeks 12 - 16)

- Range of Motion: Full, painless
- Therapeutic Exercises: Advance closed chain strengthening exercises and proprioception activities; focus on single-leg strengthening; begin elliptical
 - Straight ahead running permitted at 12 weeks
 - Swimming okay at 16 weeks
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase IV (Weeks 16 – 24): Gradual return to athletic activity

- 16 weeks: begin jumping
- 20 weeks: advance to sprinting, backward running, cutting/pivoting/changing direction
- Consider functional sports assessment

Signature:	Date:
9	