## Matthew H. Blake, MD

Team Physician, *University of Sioux Falls*Assistant Professor, Department of Orthopedic Surgery
University of South Dakota School of Medicine
www.MatthewBlakeMD.com



## Physical Therapy Prescription - Cartilage Restoration Patella / Trochlea

Name:	Date:
Procedure: R / L	Date of Surgery:
Frequency: 2-3 times per week for weeks	
PHASE I (Weeks 0 – 6): Period of protection, decrease eder	ma, activate quadriceps
extension lag  Range of Motion: Continuous Passive Motion (CPM   CPM Protocol: 1 cycle per minute starting 0  Therapeutic Exercises: Weeks 0-2: quad sets, calf pumps, passive I Weeks 2-6: PROM/AAROM to tolerance, ge hip and core exercises  Modalities: Per therapist, including electrical stimula	nproved; discontinue when able to perform SLR without 1) machine for 6-8 hours/day 0-30° (weeks 0-2), 0-60° (weeks 2-4), 0-90° (weeks 4-6) leg hangs to 45° entle patellar mobs, quad/HS/glute sets, SLR, side-lying
<ul> <li>Weightbearing: Full</li> <li>Hinged Knee Brace: None</li> <li>Range of Motion: Progress to full, painless AROM</li> <li>Therapeutic Exercises: Advance Phase I</li> <li>Modalities: Per therapist, including electrical stimula</li> </ul> Phase III (Weeks 8 – 12)	ation, ultrasound, heat (before), ice (after)
<ul> <li>Weightbearing: Full</li> <li>Range of Motion: Full, painless</li> </ul>	
Phase IV (Weeks 12 – 24)	
Advance Phase III exercises; focus on core/glutes; a	dvance to elliptical, bike, and pool as tolerated
Phase V (>6 months): Gradual return to athletic activity	
<ul> <li>Encourage maintenance program</li> <li>Return to sport-specific activity and impact when clear</li> </ul>	ared by MD at 8-9 months postop
Signature:	Date: